
Fetal bovine serum (FBS), charcoal stripped

Product Codes :

FB-1285F	FB-1001F	FB-1058F	FB-1061F	FB-1280F
FB-1345F	FB-1350F	FB-1360F	FB-1365F	FB-1003F

Collected from the source :

When searchers choose their serum an important factor that should be taken into consideration is the source, which also emphasises the traceability of the serum.

Our system of vertical integration allows us to be certain of the origins and traceability of our FBS.

Each manufactured batch is rigorously controlled, from the collection of serum and throughout all stages of its treatment and production through to final packaging on our premises.

Biosera Fetal Bovine Serum is derived from clotted whole blood aseptically collected from fetus via cardiac puncture.

The serum is imported and treated in agreement with the European regulations.

Filtration :

Final Filter Size : 0.2µm

Sterility :

All sera are tested for the absence of aerobic and anaerobic bacteria, fungi, yeast and *Mycoplasma*.

The sterility test is based on the European Pharmacopoeia requirements.

The sera are tested for the absence of *Mycoplasma* by culture.

Virus Tested :

All of our sera are tested for:

- Bovine Viral Diarrhoea (BVD)
- Cytopathogenic agents e.g. Infectious Bovine Rhinotracheitis (IBR) / BHV-1
- Hemadsorbing agents e.g. Parainfluenza Type 3 (PI3)

Sera are tested for the absence of the indicated viruses by inoculation to permissive cells. The revelation is made by immunofluorescence for pestiviruses. Cytopathogenic agents and hemadsorbing agents are detected by microscopic observations.

Endotoxin :

All sera are tested to determine the levels of endotoxins. Biosera carries out a chromokinetic quantitative test, according to the method D of the European Pharmacopoeia.

The endotoxin reagent is standardized against the US reference endotoxin.

Haemoglobin :

The haemoglobin level is measured by spectrophotometer.

Osmolality :

Determined by a lowered freezing temperature. The osmometer is calibrated against standard solutions.

Cell Culture :

Biological performance is assessed using cell culture medium supplemented with the serum being tested.

During the test period, cultures are examined microscopically for any morphological abnormalities that may indicate toxic components in the serum.

Cell Culture Tests :

Cell Growth, Plating Efficiency, Cloning Efficiency.

Cell Lines Tested :

The following cell lines are tested with the serum:

HELA -Cancer Cell/Human.

L929 -Fibroblast-Mouse/ As Macrophage

SP2/O-AG14 -Mouse/Lymphoma

MRC- 5 -Human/Lung.

Total Protein :

Determined by Biuret Colorimetry.

Treatment :

Charcoal stripped serum is treated by filtering chilled serum through an activated carbon adsorbent filter to remove non-polar material. This treatment removes lipophilic material but has little effect on the concentration of salts dissolved in the serum.

Applications :

Biosera offers Charcoal / Dextran stripped foetal bovine serum for researchers requiring low levels of various hormones. Charcoal / Dextran stripping reduces the concentration of steroid hormones in serum eg estradiol, progesterone, cortisol, testosterone, T3 and T4. It has been demonstrated and used in numerous studies both *in vivo* and *in vitro* (e.g., 1,2,3,4). So this serum is useful for utilisation in receptor studies, oestrogen related investigations, or when endogenous steroid hormones may interfere with experimental work.

In addition, Charcoal / Dextran treatment has been demonstrated to improve immunoassay systems (5,6,7,8,9,10,11) ; Herbert, et al. (12) showed that the use of Charcoal / Dextran improved insulin assay methods.

Moreover, some studies indicate that Charcoal / Dextran treatments may be used to minimise lot to lot serum variability.

This serum may show some reduced growth promotion of cells requiring the presence of certain hormones and growth factors.

The level of endotoxin is in general higher than non-treated serum because of the endotoxins in the charcoal and in the dextran.

Country of Origin :

The country in which the serum was taken from the donor/animal.

Biosera sera are sourced from the following countries

FB-1058F Uruguay

FB-1345F Central America

FB-1365F Chile

FB-1061F Dominican republic

FB-1280F France

FB-1360F Mexico

FB-1001F South America

FB-1350F USA

FB-1285F Ireland

FB-1003F South Africa

Storage conditions :

Store at -20°C

Shelf life :

5 years

Recommended use :

- Respect storage conditions of the serum
- Do not use the serum after its expiry date
- Store serum in an area protected from light
- Manipulate serum in aseptic conditions (e.g. : under laminar air flow)
- Wear clothes adapted to the manipulation of serum to avoid contamination (e.g. : gloves, mask, hygiene cap, overall...)
- In order to preserve all serum qualities, it is recommended to thaw out the flask, to aliquote, then to re-freeze the produced flasks rather than to thaw out and re-freeze the flask at each use.
- It is recommended to use the serum immediately after its thaw out. However, if it is not useful, it is possible to store thaw out serum, at +2°C / +8°C, until 26 weeks without significant decrease of its performances in cell culture.

The product is intended to be used in vitro, in laboratory only. Do not use it in therapy, human or veterinary applications.

Note:

The raw serum may be gamma irradiated before filtration for different reasons:

- Importation regulation
- Exportation necessity (all the serum intended for the Chinese market is gamma irradiated at minimum 25kGy)
- Technical or quality aspects.

To be informed if your batch is concerned by the gamma irradiation before filtration, please contact Biosera.

REFERENCES :

1. Carson, R.S., D.M. Robertson and J.K. Findlay. 1988. Ovine follicular fluid inhibits thymidine incorporation by 3T3 fibroblasts *in vitro*. J. Reprod. Fertil. 82: 447-455.
2. Coezy, E., J. Bouhnik, E. Clauser, F. Pinet, M. Philippe, J. Menard and P. Corvol. 1984. Effects of glucocorticoids and antiglucocorticoid on angiotensinogen production by hepatoma cells in culture. In Vitro 20: 528-534.
3. Friedman, E.A., M.J. Saltzman, B.G. Delano and M.M. Beyer. 1978. Reduction in hyperlipidemia in hemodialysis patients treated with charcoal and oxidized starch (oxystarch). Am. J. Clin. Nutr. 31: 1903-1914.
4. Seaver, S.S., J. van der Bosch and G. Sato. 1984. The chick oviduct in tissue culture. I. Initial characterization of growing primary oviduct tissue cultures. Exp. Cell Res. 155: 241-251.
5. Croze, F. and P. Francimont. 1984. Biological determination of inhibin in rat ovarian-cell culture medium. J. Reprod. Fertil. 72: 237-248.
6. Ding, Y.Q. and I. Huhtaniemi. 1989. Human serum LH inhibitor(s): behaviour and contribution to in vitro bioassay of LH using dispersed mouse Leydig cells. Acta Endocrinol. (Copenh.) 121: 46-54.
7. Helin, H.J., J.J. Isola, M.J. Helle and H. Adlercreutz. 1988. Influence of endocrine status on biochemical and immunocytochemical estrogen and progesterone receptor assays in breast cancer patients. Breast Cancer Res. Treat. 12: 67-73.
8. Kohno, T. and E. Ishikawa. 1986. A highly sensitive enzyme immunoassay of anti-insulin antibodies in guinea pig serum. J. Biochem. (Tokyo) 100: 1247-1251.
9. Nakamura, S., S. Sakata, T. Komaki, K. Kamikubo, K. Yasuda and K. Miura. 1986. An improved and simplified method for the detection of thyroid hormone autoantibodies (THAA) in serum. Endocrinol. Jpn. 33: 415-422.
10. Silbert, C.K. and C.T. Sawin. 1975. Double-antibody radioimmunoassay of serum insulin: effect of use of hormone-depleted human serum. Clin. Chem. 21: 1520-1522.
11. Yoshimura, M., Y. Ochi, T. Hachiya and T. Miyazaki. 1975. Estimation of the maximal T4-binding capacity of TBG using the Triosorb test in serum treated with dextran-coated charcoal. Endocrinol. Jpn. 22: 199-205.
12. Herbert V., K.-S. Lau, C.W. Gottlieb and S.J. Bleicher. 1965. Coated charcoal immunoassay of insulin. J. Clin. Endocrinology 25:1375-1384.